

## 演習問題 6 解答

問題 6.1

$$\frac{2F}{\pi d^2}$$

問題 6.2

$$\text{比ねじり角} \frac{32T}{\pi G(d_A^4 + d_1^4 - d_0^4)}$$

問題 6.3

5.35 kNm

問題 6.4

$$(1) \theta = \frac{32TL_A L_B}{\pi(G_A a^4 L_B + G_B b^4 L_A)}$$

$$(2) \frac{aG_A}{\tau_{yA} L_A} = \frac{bG_B}{\tau_{yB} L_B}$$

問題 6.5

$$(1) \frac{16FDL}{G\pi} \left( \frac{2}{d_2^4} - \frac{1}{d_1^4} \right)$$

$$(2) \tau_1 = \frac{-8FD}{\pi d_1^3}, \quad \tau_2 = \frac{8FD}{\pi d_2^3} \quad \text{丸棒 I が先に破断する}$$

問題 6.6

$$(1) T_c = 2T_0$$

$$(2) \frac{1120T_0}{\pi d^3 G}$$

問題 6.7

理由は各自で考えること

- (1) (a), (b), (c)  
 (2) (c), (a), (b)  
 (3) (a), (b), (c)  
 (4) (b), (a), (c)

問題 6.8

$$(1) \text{表面において } \tau_I = \frac{2T}{\pi a^3}, \quad \tau_{II} = 0 \quad \text{中心において } \tau_I = 0, \quad \tau_{II} = 0$$

$$(2) \theta_A = \frac{32T}{G\pi(2a)^4}16a = \frac{32T}{\pi Ga^3}$$

$$(3) \tau_I = \frac{T}{\pi a^3}, \quad \tau_{II} = -\frac{T}{\pi a^3}$$

$$(4) \theta_B = \frac{16T}{\pi Ga^3}$$

$$(5) \tau_I = \frac{DG}{32} + \frac{T}{\pi a^3}, \quad \tau_{II} = \frac{DG}{32} - \frac{T}{\pi a^3}$$

## 問題 6.9

$$(1) \tau_A = \frac{1}{4\pi R^3}T_1, \quad \tau_B = \frac{2}{\pi R^3}T_1$$

$$(2) \tau_A = \frac{2RG}{17L}D_2, \quad \tau_B = \frac{16RG}{17L}D_2$$

$$(3) \tau_A = \frac{2RG}{L}D_3, \quad \tau_B = -\frac{RG}{L}D_3$$

$$(4) \tau_A = \frac{4}{17\pi R^3}T_4, \quad \tau_B = -\frac{2}{17\pi R^3}T_4$$